

Organic **AG** Products

EX-ICUTE

Bioinsecticide/fungicide/sanitiser derived from all natural, plant produced 25-B essential oils. Aids in the control of sucking/chewing insects, mites and some fungal diseases including mildews, rust and sigatoka.



DIRECTIONS FOR USE:

Mix with adequate volumes of water before applying as a foliar spray, using a conventional knapsack sprayer. Do not apply more than one in a five day period. Coverage is essential to establish control. The use of Sticker/spreader may increase contact and efficacy of treatment. Spray early in the morning or late afternoon.

CONTROLS

Bagrada Bug (bagrada bilaris) - stink bugs & lygus
At rates of 4 oz/gal of concentrate or 1 gal/Acre in 30 gallons of water
For maintaining of Insect Control, apply at 1/2 above rate every 14 days.

NET CONTENTS - 2.5 Gal. 21lbs. / 9.6Kg at 68°F

Mites on Strawberries
Apply at rate of 2 oz/gal of water
1/2 gal/A in 30 gallons of water

Thrips on Peppers
Apply at rate of 4 oz/gal of water
or 1 gal/A in 30 gallons of water

GENERAL INFORMATION

- Use on the crop or as a post-harvest wash to kill targeted pests organically.
- Can be used for nuisance pests like houseflies or mosquitos, in and around homes, people, pets, poultry and livestock.
- Ex-icute provides farmers, growers, pest-control operators and homeowners with a novel active ingredient to prevent pesticide resistance with no residue or pre-harvest intervals.

DOSAGE

| CROP | PEST | RATE PER HA | DOSE / KNAPSACK | SPRAY INTERVAL |
|--|---|-------------|-----------------|---|
| Tomatoes, Eggplant, Pepper, Cauliflower, Watermelon, Onion, Cabbage etc. | Aphids, Thrips, Leaf Miner, Mealy bugs, Spider mites, Whitefly Armyworms. | 1.5 - 2.5lt | 100 - 250g | Begin spray when disease or pest appears and repeat every 10-14 days. |
| Mango, Pawpaw, Citrus, Avocado etc. | Whitefly, Spider mites, Scales, Fruit fly. | 1.5 - 2.5lt | 100 - 250g | Begin spray when disease or pest appears and repeat every 10-14 days. |
| Cowpeas, Soybeans, Maize, Rice, Sorghum etc | Armyworm, Stem Borer, Rust, Sogata Pyricularia oryzae, Aphids, Thrips, White fly. | 1.5 - 2.5lt | 100 - 250g | Begin spray when disease or pest appears and repeat every 10-14 days. |
| Banana | Sigatoka, Bacteriosis, Mosca Berne, White Fly. | 1.5 - 2.5lt | 100 - 250g | Begin spray when disease or pest appears and repeat every 10-14 days |
| Roses, Red Orchids, etc. | Aphids, Thrips, Leaf Miner, Mealy bugs, Spider mites. | 1.5 - 2.5lt | 100 - 250g | Twice every month. |

SAFETY PRECAUTIONS

- Avoid contact with skin, eyes.
- Keep out of reach of children.
- Do not swallow.

FIRST AID

IF IN EYES: Wash with plenty of water for at least 15 mins.
IF ON SKIN OR CLOTHING: Wash with plenty of water and soap.
IF INHALED: Move person to fresh air.

PUBLIC HEALTH RECOMMENDATION

For the control of insects, pests including mosquitos, houseflies etc in farm, houses, lawn use, 1% strength solution.

NOTICE TO THE BUYER:

Sellers guarantee is limited to the terms set out on the label and subject thereto. The buyer assumes the risk to persons or property arising from the use and handling of this product and accepts the product on that condition.

OrganicAGProducts.com

FOR SALES CONTACT: AGMERICA
INFO@AGMERICA.US

FOR AGRICULTURAL
USE ONLY.



**SHAKE WELL
BEFORE USING**

2 PER CASE WT. 44lbs. / 18.2kg @ 68°F